

## ORIGINAL ARTICLE

## Recovery of insulin sensitivity and optimal body composition after rapid weight loss in obese dogs fed a high-protein medium-carbohydrate diet

A. André<sup>1</sup>, I. Leriche<sup>2</sup>, G. Chaix<sup>3</sup>, C. Thorin<sup>1</sup>, M. Burger<sup>1</sup> and P. Nguyen<sup>1</sup>

<sup>1</sup> UBL, Oniris, National College of Veterinary Medicine, Food Science and Engineering Nantes-Atlantic, Nutrition, Physiopathology and Pharmacology Unit (NP3), Nantes, France

<sup>2</sup> Virbac Nutrition, Vauvert, France, and

<sup>3</sup> Virbac Medical Department, Carros, France

### Summary

This study investigated the effects of an experimental high-protein medium-carbohydrate diet (protein level, 46% metabolizable energy, ME). First, postprandial plasma glucose and insulin kinetics were determined in steady-state overweight/obese Beagle dogs (28%–41% excess body weight) for an experimental high-protein medium-carbohydrate diet (protein level, 46% ME) and a commercial high-carbohydrate medium-protein diet (protein level, 24%ME) in obese dogs. Secondly, all the dogs were included in a weight loss programme. They were fed the high-protein medium-carbohydrate diet, and the energy allocation was gradually reduced until they reached their optimal body weight. Insulin sensitivity and body composition were evaluated before and after weight loss using a euglycaemic–hyperinsulinaemic clamp and the deuterium oxide dilution technique respectively. For statistical analysis, linear mixed effect models were used with a significance level of 5%. Postprandial plasma glucose and insulin concentrations were substantially lower with the high-protein medium-carbohydrate diet than the high-carbohydrate medium-protein diet. These differences can be explained mainly by the difference in carbohydrate content between the two diets. Energy restriction (35% lower energy intake than in the obese state) resulted in a  $2.23 \pm 0.05\%$  loss in body weight/week, and the dogs reached their optimal body weight in 12–16 weeks. Weight loss was associated with a significant increase in insulin sensitivity. The high-protein medium-carbohydrate diet allowed fat-free mass preservation despite a relatively high rate of weekly weight loss. The increase in insulin sensitivity indicated improved control of carbohydrate metabolism, possible due to weight loss and to the nature of the diet. Thus, a high-protein medium-carbohydrate diet is a good nutritional solution for managing the weight of overweight dogs. This diet may improve glycaemic control, which could be beneficial for preventing or managing impaired glucose tolerance in obese dogs and for safe and successful weight loss while preserving lean body mass.

**Keywords** overweight, starch, glucose, metabolism, clamp, lean body mass

**Correspondence** P. Nguyen, Nutrition, Pharmacology and Physiopathology Unit, Oniris - National Veterinary School of Nantes, BP 40307, Route de Gachet, 44307 Nantes Cedex 03. Tel: +33240687635; Fax: +33240687746; E-mail: patrick.nguyen@oniris-nantes.fr

Received: 25 January 2017; accepted: 6 April 2017

### Introduction

Surveys of obesity in the canine population that have been performed worldwide indicate that the prevalence of canine obesity has been increasing for years (Colliard et al., 2006; Laflamme, 2006; Courcier et al., 2010; Mao et al., 2013). Obese dogs are those with a body condition score of 4/5 or 5/5 according to the WSAVA Nutritional Assessment Guidelines (WSAVA Nutritional Assessment Guidelines Task Force Members, 2011). According to the

Association for Pet Obesity Prevention (<http://www.petobesityprevention.org>), as many as 53.8% of US dogs would have been overweight or obese in 2015. Canine obesity is associated with important metabolic and hormonal changes (Gayet et al., 2004; Clark and Hoenig, 2016), such as impaired glucose tolerance, reduced sensitivity to insulin (German et al., 2010b; Laflamme, 2011) and dyslipidaemia (Baillhache et al., 2003; Verkest et al., 2012). Moreover, obesity is considered a significant risk factor in the development of chronic diseases, such as renal

disease, heart disease and cancer (Thengchaisri *et al.*, 2014; Weeth, 2016).

With respect to glucose intolerance and reduced insulin sensitivity, approaches that use low glycaemic index diets have been tested to improve weight and metabolic outcomes (Mitsuhashi *et al.*, 2012). Foods with low glycaemic index may also reduce oxidative stress and protect the cardiovascular system in dogs (Adolphe *et al.*, 2012). The glycaemic index was developed to rank carbohydrate-containing foods according to their effects on blood glucose levels in humans (Jenkins *et al.*, 1981). Rapidly available glucose (RAG) is defined as the amount of glucose released *in vitro* from a food during the first 20 min of incubation with adequate enzymes, and it has been shown to be related to the glycaemic response in human studies (Englyst *et al.*, 1999). In rats, both plasma glucose and insulin responses were positively correlated to the rate of hydrolysis with  $\alpha$ -amylase *in vitro* (Holm *et al.*, 1988). However, a more recent human study showed that two carbohydrate sources with similar RAG content elicited similar glycaemic responses but different insulin responses (Al-Mssallem *et al.*, 2011). Thus, predicting the effects of a diet on postprandial plasma glucose and insulin kinetics is not easy, and field trials in target species are necessary to accurately evaluate the *in vivo* responses.

In dogs, postprandial glucose kinetics are mainly influenced by the amount of starch in the diet (Nguyen *et al.*, 1994). Nevertheless, many factors other than available carbohydrate level affect postprandial glycaemia and insulinaemia, including starch source and structure, macronutrient content and source (Kendall *et al.*, 2006). In particular, it has been shown, in humans, that high protein intake increases insulin secretion (Gulliford *et al.*, 1989), and moreover that different sources of protein may lead to different insulin responses (Nuttall and Gannon, 1990).

Compared to the starch level, the protein level has a limited impact on postprandial glycaemia, but there has long been interest in the use of high-protein diets for weight loss. Orthopaedic or other surgical procedures may require weight loss, either for causal reasons (orthopaedic procedures) or because obesity makes anaesthetic management more difficult or is a risk factor for surgical site infections (Love and Cline, 2015). Therefore, high-energy restriction may be chosen when there is a medical indication for rapid weight loss. In dogs, energy restriction is suggested for losses of 1%–2% of the initial body weight (BW) per week (Nguyen and Diez, 2010). In all cases, it takes months to lose the excess 25%–30% of excess BW that is, the weight over the optimal BW. Although the

literature is not entirely conclusive on this point, in general, the risk of losing lean body mass is higher with a higher rate of weight loss. In such cases, attention should be paid to the composition of the diet and to the provision of essential nutrients, including protein. Lower lean body mass loss was observed during weight loss when dogs were given a high-protein diet (crude protein, CP: 39% metabolizable energy, ME) compared to a medium-protein diet (CP: 20%ME) (Hannah, 1999). In a previous study we conducted in obese dogs, we showed that a high-protein diet (CP: 41%ME) allowed the conservation of lean body mass despite a high rate of weight loss (2.6% loss per week) (Blanchard *et al.*, 2004). More recently, two other high-protein diets (CP: ~40%ME) were also shown to allow safe weight loss in dogs, but with a lower rate of weight loss of 0.7%–1.0% per week (German *et al.*, 2010a).

In the present study, we compared two diets with protein levels of 46%ME or 24%ME. Although the terms do not have widely accepted definitions, these diets can be considered high-protein and medium-protein diets, respectively, in accordance with terminology used in previous studies and in accordance with the minimum FEDIAF recommendation (2016) for maintenance (CP: 18%ME), which is the lowest acceptable level.

The objectives of our research were twofold. Study 1 compared the postprandial glycaemic and insulinaemic kinetics induced by a high-protein medium-carbohydrate (HP-MC) diet vs. a high-carbohydrate medium-protein (HC-MP) diet in overweight/obese dogs. Study 2 assessed the effects of the HP-MC diet on body composition and insulin sensitivity in overweight/obese dogs, energy restricted to induce rapid weight loss.

## Materials and methods

### Animals

Nine adult overweight/obese Beagle dogs (five neutered males and four spayed females) aged  $3.2 \pm 0.4$  years were included in these studies. The experimental protocols adhered to European Union guidelines and followed the requirements for animal welfare of the French Ministry of Agriculture. This work was approved by the Animal Use and Care Advisory (Comité d'Éthique en Expérimentation Animale des Pays de la Loire; approval number 00281.02). The dogs were housed individually in an air-conditioned environment. The room temperature was maintained between 19 °C and 24 °C. The dogs were housed in the same room with permanent visual contact with

each other to prevent isolation stress. As a group, they went outdoors daily to interact and exercise for a limited and controlled period of time. Water was freely available at all times.

Before the two studies began, dogs with an optimal BW of  $13.1 \pm 0.7$  kg were fed a high-energy diet and developed obesity. Given the same diet at  $\sim 150$  kcal (628 kJ) ME/kg optimal BW<sup>0.75</sup>, the dogs were maintained in an overweight or obese state for 30 weeks.

At the beginning of Study 2, the dogs' mean BW was  $17.6 \pm 0.9$  kg. The mean percentage of excess BW was 34% (range: 28%–41%), and the dogs' body condition score (BCS) was 7 ( $n = 1$ ) or 8 ( $n = 8$ ) on a nine-point scale (Laflamme, 1997). BW and BCS were monitored and recorded weekly.

## Diets

Two diets were used in Study 1: a commercial HC-MP medium-energy diet, and a nutritionally balanced experimental HP-MC low-energy diet that was not commercially available at the time of the study. The

**Table 1** Nutritional characteristics of an experimental high-protein medium-carbohydrate (HP-MC) diet and a commercial high-carbohydrate medium-protein (HC-MP) diet: chemical analysis (as fed) and contribution of macronutrients to metabolizable energy (%ME). The two diets were both used in an acute feeding trial (Study 1), and the HP-MC diet was also used in a long-term feeding trial (Study 2)

|                                    | HP-MC diet | HC-MP diet |
|------------------------------------|------------|------------|
| Moisture (%)                       | 7.0        | 7.4        |
| Crude protein (%)                  | 37.5       | 22.4       |
| Crude fat (%)                      | 11.4       | 13.5       |
| Nitrogen-free extract (%)          | 23.0       | 41.6       |
| Crude fibre (%)                    | 13.8       | 10.8       |
| Starch (%)                         | 13.4       | 31.9       |
| Total sugars (%)                   | 0.7        | 0.3        |
| Insoluble fibre (%)                | 22.8       | 17.0       |
| Soluble fibre (%)                  | 1.6        | 0.8        |
| Ash (%)                            | 7.3        | 4.3        |
| Metabolizable energy* (kcal/100 g) | 293        | 332        |
| Metabolizable energy* (kJ/100 g)   | 1230       | 1390       |
| Protein (%ME)*                     | 46         | 23         |
| Starch (%ME)*                      | 19         | 41         |
| Fat (%ME)*                         | 31         | 36         |

\**in vivo* determination.

HP-MC diet contained dehydrated pork and poultry proteins, lignocellulose, potato starch, hydrolysed animal protein, animal fat, field bean hulls, minerals, dried beet pulp, psyllium fibre, fructo-oligosaccharides, chitosan, chondroitin sulphate, pasteurized *Lactobacillus acidophilus*, vitamins.

HC-MP contained maize, dehydrated poultry proteins, rice, cellulose, animal fat, hydrolysed animal protein, pea fibre, dried whole egg, vegetable oil, linseed, minerals, vitamins.

chemical composition of each diet is presented in Table 1. In Study 2, only the HP-MC diet was used.

## *In vitro* measurement of RAG in the diets

The measurement of RAG in the two diets was performed by Agrobio (Vezin Le Coquet, France). The *in vitro* technique (Englyst et al., 1999) is based on HPLC measurement of the glucose released from the food after a 20-min incubation with digestive enzymes (pepsin, pancreatin, invertase and amyloglucosidase).

## Study design

In Study 1, eight dogs, in stable overweight or obese state, were subjected, in a crossover design, to postprandial plasma glucose and insulin kinetics assays while they were fed either the HP-MC diet or the HC-MP diet. Between the two kinetics assays, the dogs were fed, for 1 week, the diet used for the development of obesity. In Study 2, nine dogs were subjected to a weight loss programme with the HP-MC diet. The target was the recovery of optimal BW. The dogs were fed once daily. The initial food allocation was 120 kcal (503 kJ) ME/kg optimal BW<sup>0.75</sup>, which represents an approximate 20% decrease relative to the previous diet. The target weekly weight loss rate was 2%–2.5% (compared to the BW of the previous week). The food allowance was decreased by 10% as many times as necessary when an insufficient change in BW was observed over a 2-week period. The duration of the study was not predetermined; it lasted until each dog reached its optimal BW. Body composition and insulin sensitivity were evaluated before the weight loss intervention and when the target weight was achieved (see below for details).

## Postprandial kinetics and biochemical analyses (Study 1)

On the morning of the plasma glucose and insulin kinetics assays (i.e., the test days) and after a 24-h food deprivation period, eight overweight/obese dogs were fed either the HP-MC or HC-MP diet in a single meal. This comprised 544 kJ/130 kcal ME/kg optimal BW<sup>0.75</sup>, which was the nearest value to their consumption in the stable obese state, and was in the range of FEDIAF recommendations (2016) for BW maintenance in adult dogs. The postprandial kinetics assays were performed using a crossover design in that during the week between the two test days, the dogs were fed the diet that was used for the development of obesity. All dogs consumed their meal in less than

10 min. Jugular vein blood samples were collected before the meal, immediately after the meal, and then 10, 20, 30, 45, 60, 90, 120, 150, 180, 210, 240, 300 and 360 min after the end of the meal. Blood glucose concentrations were determined in whole blood using a portable blood glucose meter (Alphatrak, Abbott Laboratories, IL, USA) that was validated for use in dogs (Kang et al., 2015). The heparinized blood samples were immediately refrigerated on ice and centrifuged ( $2200 \times g$ , 10 min, 4 °C). The plasma was collected and stored at  $-20$  °C until analysis. Plasma insulin was determined by radioimmunoassay (Insulin IRMA kit, Beckman Coulter, Prague, Czech Republic).

#### Euglycaemic–hyperinsulinaemic clamp (Study 2)

A 3-h euglycaemic–hyperinsulinaemic clamp was used to estimate insulin sensitivity before and after weight loss with the HP-MC diet in seven 24-h unfed dogs as described previously (Bailhache et al., 2003).

#### Body composition (Study 2)

Body composition was determined three times: before the period of weight gain, in the stable overweight/obese state and after the weight loss programme. Total body water (TBW) was assessed using an isotopic dilution of  $^2\text{H}_2\text{O}$ . After a 24-h food deprivation period, dogs were confined to metabolism cages. Water was withheld, allowing body water equilibration from 2 h before to 4 h after the tracer injection. Dogs were injected subcutaneously (0.50 g/kg BW) with physiological saline (9 g NaCl/L) prepared with  $^2\text{H}_2\text{O}$  (99.9%  $^2\text{H}/\text{H}$ , Eurisotop, Gif-sur-Yvette, France). Venous blood samples were obtained prior to and 4 h after injection of the tracer. EDTA plasma was stored at  $-20$  °C in sealed vials until analysis. Deuterium enrichment ( $^2\text{H}/\text{H}$ ) in plasma water was determined by Fourier transform infrared spectroscopy on a Vector 22-type spectroscope (Brüker SA, Wissembourg, France) (Ferrier et al., 2002). TBW was determined from the dilution space of the isotope. Fat-free mass (FFM) was calculated as the TBW/hydration rate of FFM using a canine-specific hydration rate of 0.744 (Harrison et al., 1936).

#### Data analysis

All data were expressed as means  $\pm$  standard deviation. To take into account the crossover design of the postprandial kinetics study, the two diets were compared using a linear mixed effects model analysis and using Tukey's tests. For an accurate analysis

of the repeated measures design, a mixed effects model was used for body composition and insulin sensitivity index comparisons. Analyses were performed in R software (R Core Team, 2013) using the nlme and multcomp packages. The nlme package that was provided by the Rcore team contains all of the functions used for analysis in a mixed effects model (linear or nonlinear). We considered  $p < 0.05$  to indicate a significant difference between data sets.

## Results

### Study 1

#### *Postprandial glycaemic and insulinaemic kinetics*

The kinetics of the glucose (Fig. 1a) and insulin (Fig. 1b) responses were different between the two tested diets. The glucose peak appeared earlier for the HP-MC diet than for the HC-MP diet. The maximum glucose increase did not differ significantly, but the area under the glucose response curve was approximately double for the HC-MP diet compared to the HP-MC diet ( $44 \pm 35$  g.min/l vs.  $21 \pm 10$  g.min/l,  $p = 0.044$ ). The maximum plasma insulin response was significantly lower for the HP-MC diet than for the HC-MP diet ( $32 \pm 11$   $\mu\text{U}/\text{ml}$  vs.  $68 \pm 39$   $\mu\text{U}/\text{ml}$ ,  $p = 0.031$ , Table 2). The incremental area under the insulin response curve, which could be considered a first approximation of the total insulin secretion, was approximately threefold higher for the HC-MP diet than for the HP-MC diet ( $12\,400 \pm 6400$   $\mu\text{U}\cdot\text{min}/\text{ml}$  vs.  $3600 \pm 2500$   $\mu\text{U}\cdot\text{min}/\text{ml}$ ,  $p = 0.004$ ). The average postprandial insulinaemia was significantly higher for the HC-MP diet ( $47.1 \pm 13.2$   $\mu\text{U}/\text{ml}$  vs.  $30.3 \pm 9.3$   $\mu\text{U}/\text{ml}$ ,  $p = 0.002$ ).


#### *Determination of RAG in foods*

The estimated values of the *in vitro* glycaemic index were 20 and 44 (on a scale of 0–100) for the HP-MC and HC-MP diets respectively.

### Study 2

#### *BW, BCS and characteristics of the weight loss period*

Dogs reached their target weight, which corresponded to their optimal BW ( $13.1 \pm 0.7$  kg and a BCS of 5), after 12–16 weeks (mean,  $14.3 \pm 1.4$  weeks). The mean weight loss rate was  $2.23 \pm 0.05\%$  per week. The mean daily energy allowance was  $98 \pm 3$  kcal ( $410 \pm 13$  kJ) ME/optimal BW<sup>0.75</sup> over the entire weight loss period. The mean daily consumption of protein was  $12.9 \pm 2.5$  g/kg optimal BW<sup>0.75</sup>.



**Fig. 1** Postprandial incremental changes in plasma glucose concentration (a) and insulin concentration (b) from time 0 values in obese dogs fed a high-protein medium-carbohydrate (HP-MC) diet or a high-carbohydrate medium-protein (HC-MP) diet (Study 1;  $n = 8$ ). Values are expressed as means.


**Table 2** Plasma glucose and insulin response characteristics (mean, maximum increments and area under the curves) in obese dogs after a meal-feeding test using a high-protein medium-carbohydrate (HP-MC) diet or a high-carbohydrate medium-protein (HC-MP) diet (Study 1;  $n = 8$ ). Values are expressed as means  $\pm$  SD

|  | HP-MC diet      | HC-MP diet        | p  |
|--|-----------------|-------------------|----|
| Mean postprandial glycaemia (g/l)            | 1.05 $\pm$ 0.10 | 1.11 $\pm$ 0.13   | ns |
| Mean postprandial insulinaemia ( $\mu$ U/ml) | 30.3 $\pm$ 9.3  | 47.1 $\pm$ 13.2   | ** |
| AUCG (g $\times$ min/l)                      | 21 $\pm$ 10     | 44 $\pm$ 35       | *  |
| AUCI ( $\mu$ U $\times$ min/ml)              | 3600 $\pm$ 2500 | 12 400 $\pm$ 6700 | ** |
| Maximum glucose increment (g/l)              | 0.23 $\pm$ 0.08 | 0.27 $\pm$ 0.16   | ns |
| Maximum insulin increment ( $\mu$ U/ml)      | 32 $\pm$ 11     | 68 $\pm$ 39       | *  |

AUCG, area under the curve for glucose; AUCI, area under the curve for insulin. \* $p < 0.05$ , \*\* $p < 0.01$ , ns no significant difference between the two tested diets. Values are expressed as mean  $\pm$  SD, where  $n = 8$  dogs.


#### Insulin sensitivity

The plasma insulin and glucose concentrations in the unfed state were not significantly different in the



**Fig. 2** Insulin sensitivity as measured in Study 2 by the euglycaemic-hyperinsulinaemic clamp method in seven dogs before and after weight loss with a high-protein medium-carbohydrate (HP-MC) diet. \*\* $p < 0.01$  between lean and obese groups. Means  $\pm$  SD are indicated.

obese dogs than in the dogs at the end of the weight loss period (data not shown). As Fig. 2 illustrates, insulin sensitivity was higher after weight loss ( $0.24 \pm 0.05$  vs.  $0.15 \pm 0.06$ ,  $p = 0.006$ ). The euglycaemic-hyperinsulinaemic clamp failed in two dogs,



**Fig. 3** Body composition of the dogs as determined in Study 2 using deuterium oxide dilution to assess (a) the fat mass (FM) as a percentage of body weight (BW) (%FM) and (b) the fat-free mass (FFM) in kg, at an optimal time (in the lean state before weight gain), in the obese state and after weight loss with a high-protein medium-carbohydrate (HP-MC) diet. Means  $\pm$  SD are indicated;  $n = 9$ . Means with different supercripts differ ( $p < 0.0001$ ).

showing unexpected and unexplained changes in plasma glucose despite constant insulin and glucose infusion rates.

#### Body composition

The body composition results are presented in Fig. 3. Weight loss resulted in a significant decrease in fat mass (FM) and FFM ( $p < 0.0001$ ). The body composition after weight loss was similar to the body composition assessed at the optimal BW.

#### Discussion

The first aim of the study was to compare the effects of two types of meals, that is, the HP-MC and HC-MP diets, on postprandial glycaemic and insulinaemic

responses in overweight/obese dogs. Starch content, which was between 0.4 and 52.7 g/100 g dry matter in the tested diets, is the primary determinant of postprandial glucose in healthy dogs (Nguyen et al., 1998). Elliott et al. (2012) showed that a restricted carbohydrate diet facilitated the control of blood glucose levels in healthy dogs: a medium-carbohydrate (25%ME) high-protein (43%ME) diet resulted in lower peak and postprandial glucose concentrations compared to a common adult maintenance diet (high carbohydrate, 55%ME; medium protein, 22%ME). Despite different levels of carbohydrates in the two tested diets in Study 1 (Starch: 19%ME vs. 40%ME) and different *in vitro* RAG concentrations (20 for the HP-MC diet vs. 44 for the HC-MP diet), we observed no significant differences in the mean postprandial plasma glucose concentration or in the maximum glucose increase. However, the HP-MC diet led to significant differences in insulinaemic responses (lower maximum increment, lower mean, lower AUCI). Our results are in agreement with those of Hewson-Hughes et al. (2011), who reported that there were minimal changes in plasma glucose concentration whatever the level of starch in the diet and that the increase in insulin concentration was lower in healthy dogs fed a low-starch diet (9.5%) compared to a medium-starch (23%) or high-starch (31.7%) diet. We hypothesize that the results we obtained on the test days would be the same as those resulting from usual and daily consumption.

There is currently no standardized method for evaluating the glycaemic index in non-human species. It would have been interesting to use the same method as Adolphe et al. (2012). For a given food that has 10 g of available carbohydrate, the glycaemic index was calculated by expressing the area under the curve for glucose as a percentage of the mean response to the glucose control (a 20% glucose solution providing 10 g of glucose) that was consumed by the same dog. In our study, the two diets were compared without normalizing the quantity of digestible carbohydrates.

Dietary fibre, another macronutrient, has been evaluated for its potential benefits in glucose supply regulation in dogs. Variations in fibre intake from 3.2 to 39.1 g of total dietary fibre/100 g DM do not significantly influence postprandial glucose and insulin responses in healthy dogs (Nguyen et al., 1998). In our investigation, although the sources and levels of fibre were different, the different levels of starch rather than the quantity and quality of the fibre might explain the results.

The question of whether high postprandial insulinaemia and glycaemia have deleterious impacts in

obese dogs still has no clear answer. Adolphe *et al.* (2014) suggested that a diet that would significantly reduce the postprandial insulin response, even without weight loss, might improve the metabolic profile of dogs. With regard to postprandial responses, the HP-MC diet appears to be more appropriate for overweight dogs. However, obesity is associated with numerous diseases (cancer, heart disease, chronic kidney disease), and the healthiest solution may be weight loss.

The second objective of this study was to investigate the effect of a diet with high-protein (46%ME) and medium-carbohydrate content (19%ME) during weight loss on FFM preservation. This diet was optimized for the nutritional management of reduction of excessive BW in obese dogs, which were thus the target animals.

The first determinant of weight lost is energy restriction: energy intake must be lower than energy expenditure. The dogs' energy consumption during the weight loss period ( $98 \pm 3$  kcal ( $410 \pm 13$  kJ) ME/kg BW<sup>0.75</sup>) may appear high, as it is in the range of the FEDIAF recommendations (2016) for maintenance in adult dogs (95–130 kcal (397–544 kJ) ME/kg BW<sup>0.75</sup>, depending on sexual status and physical activity). However, the animals included in this study were kennel dogs, and their usual food consumption provided 150 kcal (628 kJ) ME/kg BW<sup>0.75</sup>, as determined to maintain their optimal BW. The diet used to fight obesity has to be specifically balanced to cover all nutritional requirements while preserving lean mass. In the present study, dogs fed a high-protein diet quickly recovered their optimal weight, and their body composition was similar to before they gained weight. The higher absolute FFM in the obese state may be related to the increase in the mass of some organs and the mass of body fluids, as reported in humans (Bosy-Westphal *et al.*, 2009; Pourhassan *et al.*, 2014).

Although the term "high-protein diet" is used, it should be kept in mind that dogs require an absolute protein provision rather than a high dietary protein level and that the protein requirement of dogs remains a grey zone, even for a maintenance diet (minimal vs. optimal, National Research Council (2006) vs. FEDIAF (The European Pet Food Industry Federation) (2016) recommendations). In our study, the mean daily consumption of protein was  $12.9 \pm 2.5$  g/kg optimal BW<sup>0.75</sup>. The efficacy of a high-protein, low-carbohydrate, low-energy diet (CP 47.5%DM, i.e., approximately 63%ME, starch 5.3% DM, 11.6 kJ/g as fed) for weight loss has been shown in obese Beagles, as this diet allowed rapid and safe

weight loss without altering the lean mass (Diez *et al.*, 2002). The mean daily consumption of protein in that study was 15.4 g/kg optimal BW<sup>0.75</sup>. Blanchard *et al.* (2004) showed the same results with another high-protein, medium-carbohydrate, low-energy diet (CP 40%ME, carbohydrate 30%ME, 10.5 kJ/g as fed), and the mean daily consumption of protein was as low as 6.3 g/kg optimal BW<sup>0.75</sup>. In the study by Hannah (1999), a high-protein diet (CP 39%ME) resulted in a lower loss of lean tissue than a medium-protein one (CP 20%ME), while the loss of lean mass with an intermediary diet (CP: 30%ME) was not significantly different from that determined by the two other diets. Unfortunately, this study did not state how much protein the dogs consumed nor could that be calculated.

Taken together, these data may look somewhat inconsistent, but this is probably not the case. Indeed, like our study, these studies aimed to show that diets that are formulated for weight loss programmes must have a higher protein-to-energy ratio than maintenance diets in order to restrict the energy allowance while ensuring that the dogs' protein requirement is met. These studies may also have intended to show that dogs receiving an energy-restricted diet that subsequently lose weight could benefit from a higher protein provision, as was recently shown in older people for whom the protein provision (in g/kg) should be twice as high as the maintenance requirement (Weijs and Wolfe, 2016). This would not probably be the case in dogs, at least in the mentioned studies. Indeed, if a minimal protein provision is needed to counteract protein turnover and thereby ensure the preservation of lean body mass, once this goal is reached, there is probably nothing more to expect from a higher protein allowance. There are a few ways to explain differences between the dog studies and the cited human studies. First, the human studies were conducted in older people in whom relative anabolic resistance has been described (Moore *et al.*, 2015). Second, the usual daily protein allowance in dogs is already higher than their strict protein requirement. Indeed, the usual allowance is approximately three times (2.5 vs. 0.83 g/kg) that of humans, and about three times (~5 vs. 1.8 g/kg BW<sup>0.75</sup>) the minimum determined in dog nitrogen balance studies (Kendall *et al.*, 1982). In these respects, the allowance (6.3 g/kg optimal BW<sup>0.75</sup>) in the experiment in the Blanchard *et al.* (2004) study was probably high enough to ensure preservation of lean mass (compared to the previous lean state), and it is not surprising that higher protein provisions gave a similar result. Accordingly, if the dietary protein level must be higher in diets intended for weight loss than in maintenance diets (according

to FEDIAF recommendation (2016)), there is no evidence to substantiate the need for an absolute higher protein provision during weight loss than during maintenance in dogs, given that the usual recommendation is already higher than the strict requirement.

A high-protein diet is not only efficient in terms of preserving FFM, but it is also in keeping with some dogs' carnivorous nutritional traits. A study that investigated macronutrient selection in domestic dogs showed that when dogs had a choice of foods with variable protein, fat and carbohydrate content, they chose a diet with approximately one-third of the energy derived from protein, 7% from carbohydrates and more than 60% of energy from fat (Hewson-Hughes et al., 2013). Wolves, which are pack-hunting progenitors of the domestic dog, are adaptive and true carnivores. Their protein requirement is much higher than that of dogs (Bosch et al., 2015). Even though dogs have adapted to a starch-rich diet during domestication, they still share numerous traits with true carnivores (Bosch et al., 2015). A high-protein diet may more closely reflect dogs' ancestral dietary pattern. Moreover, rapid feeding, which is an adaptation to scavenging, is also a trait in this dietary pattern, and dogs that retain this tendency can rapidly become obese (Bradshaw, 2006). As humans regulate the intake of protein energy (PE) more strongly than non-protein energy (nPE), it has been proposed that the rise in obesity may have resulted partly from a shift towards diets with reduced PE:nPE ratios (Raubenheimer et al., 2015). High-protein diets could therefore be beneficial in dogs in terms of the prevention of

overweight and, at the same time, be in accordance with the ancestral dietary pattern of the dogs.

Lastly, the insulin sensitivity index showed that obese dogs are more insulin-sensitive after they lose weight. Our results showed that the changes in glucose metabolism observed in dogs that were suffering from primary obesity due to excess food intake are reversible. This improvement may be due mainly to weight loss *per se*, as demonstrated previously (German et al., 2009). Another study that used a control diet during weight loss could clarify the relative importance of weight loss and diet composition on the improvement of insulin sensitivity.

Our study showed that a HP-MC diet may be a good nutritional solution for safe weight loss in overweight/obese dogs, as it allowed rapid weight loss with preservation of lean body mass. Treating obesity could improve insulin sensitivity. A diet with a protein-to-carbohydrate ratio of 1.6 and a protein-to-energy ratio of 112 g MP/Mcal ME (28 g MP/MJ ME) was associated with lower postprandial glucose and insulin concentrations than a diet with a protein-to-carbohydrate ratio of 0.5 and a protein-to-energy ratio of 61 g MP/Mcal ME (15 g MP/MJ ME). Such a HP-MC diet may help in glycaemic control, which could be advantageous for the prevention or management of impaired glucose tolerance or diabetes mellitus in obese dogs.

### Acknowledgements

The authors thank Philippe Bleis and Samuel Ninet for their excellent technical assistance.

### References

- Adolphe, J. L.; Drew, M. D.; Huang, Q.; Silver, T. I.; Weber, L. P., 2012: Postprandial impairment of flow-mediated dilation and elevated methylglyoxal after simple but not complex carbohydrate consumption in dogs. *Nutrition Research* **32**, 278–284.
- Adolphe, J. L.; Drew, M. D.; Silver, T. I.; Foughse, J.; Childs, H.; Weber, L. P., 2014: Effect of an extruded pea or rice diet on postprandial insulin and cardiovascular responses in dogs. *Journal of Animal Physiology and Animal Nutrition* **99**, 767–776.
- Al-Mssallem, M. Q.; Hampton, S. M.; Frost, G. S.; Brown, J. E., 2011: A study of Hassawi rice (*Oryza sativa* L.) in terms of its carbohydrate hydrolysis (in vitro) and glycaemic and insulinaemic indices (in vivo). *European Journal of Clinical Nutrition* **65**, 627–634.
- Bailhache, E.; Ouguerram, K.; Gayet, C.; Krempf, M.; Siliart, B.; Magot, T.; Nguyen, P., 2003: An insulin-resistant hypertriglyceridaemic normotensive obese dog model: assessment of insulin resistance by the euglycaemic hyperinsulinaemic clamp in combination with the stable isotope technique. *Journal of Animal Physiology and Animal Nutrition* **87**, 86–95.
- Blanchard, G.; Nguyen, P.; Gayet, C.; Leriche, I.; Siliart, B.; Paragon, B. M., 2004: Rapid weight loss with a high-protein low-energy diet allows the recovery of ideal body composition and insulin sensitivity in obese dogs. *Journal of Nutrition* **134**, 2148S–2150S.
- Bosch, G.; Hagen-Plantinga, E. A.; Hendriks, W. H., 2015: Dietary nutrient profiles of wild wolves: insights for optimal dog nutrition? *British Journal of Nutrition* **113**(Suppl), S40–S54.
- Bosy-Westphal, A.; Kossel, E.; Goele, K.; Later, W.; Hitze, B.; Settler, U.; Heller, M.; Glüer, C. C.; Heymsfield, S. B.; Müller, M. J., 2009: Contribution of individual organ mass loss to weight loss-associated decline in resting energy expenditure. *American Journal of Clinical Nutrition* **90**, 993–1001.
- Bradshaw, J. W., 2006: The evolutionary basis for the feeding behavior of domestic dogs (*Canis familiaris*) and cats (*Felis catus*). *Journal of Nutrition* **136**, 1927S–1931S.
- Clark, M.; Hoenig, M., 2016: Metabolic effects of obesity and its interaction with endocrine diseases. *Veterinary Clinics of North America: Small Animal Practice* **46**, 797–815.
- Colliard, L.; Ancel, J.; Benet, J. J.; Paragon, B. M.; Blanchard, G., 2006: Risk



- factors for obesity in dogs in France. *Journal of Nutrition* **136**, 1951S–1954S.
- Courcier, E. A.; Thomson, R. M.; Mellor, D. J.; Yam, P. S., 2010: An epidemiological study of environmental factors associated with canine obesity. *Journal of Small Animal Practice* **51**, 362–367.
- Diez, M.; Nguyen, P.; Jeusette, I.; Devois, C.; Istasse, L.; Biourge, V., 2002: Weight loss in obese dogs: evaluation of a high-protein, low-carbohydrate diet. *Journal of Nutrition* **132**, 1685S–1687S.
- Elliott, K. F.; Rand, J. S.; Fleeman, L. M.; Morton, J. M.; Litster, A. L.; Biourge, V. C.; Markwell, P. J., 2012: A diet lower in digestible carbohydrate results in lower postprandial glucose concentrations compared with a traditional canine diabetes diet and an adult maintenance diet in healthy dogs. *Research in Veterinary Science* **93**, 288–295.
- Englyst, K. N.; Englyst, H. N.; Hudson, G. J.; Cole, T. J.; Cummings, J. H., 1999: Rapidly available glucose in foods: an in vitro measurement that reflects the glycemic response. *American Journal of Clinical Nutrition* **69**, 448–454.
- FEDIAF (The European Pet Food Industry Federation) 2016: *Nutritional Guidelines for Complete and Complementary Pet Food for Cats and Dogs*.
- Ferrier, L.; Robert, P.; Dumon, H.; Martin, L.; Nguyen, P., 2002: Evaluation of body composition in dogs by isotopic dilution using a low-cost technique. Fourier-transform infrared spectroscopy. *Journal of Nutrition* **132**, 1725S–1727S.
- Gayet, C.; Bailhache, E.; Dumon, H.; Martin, L.; Siliart, B.; Nguyen, P., 2004: Insulin resistance and changes in plasma concentration of TNF $\alpha$ , IGF1, and NEFA in dogs during weight gain and obesity. *Journal of Animal Physiology and Animal Nutrition* **88**, 157–165.
- German, A. J.; Hervera, M.; Hunter, L.; Holden, S. L.; Morris, P. J.; Biourge, V.; Trayhurn, P., 2009: Improvement in insulin resistance and reduction in plasma inflammatory adipokines after weight loss in obese dogs. *Domestic Animal Endocrinology* **37**, 214–226.
- German, A. J.; Holden, S. L.; Bissot, T.; Morris, P. J.; Biourge, V., 2010a: A high protein high fibre diet improves weight loss in obese dogs. *Veterinary Journal* **183**, 294–297.
- German, A. J.; Ryan, V. H.; German, A. C.; Wood, I. S.; Trayhurn, P., 2010b: Obesity, its associated disorders and the role of inflammatory adipokines in companion animals. *Veterinary Journal* **185**, 4–9.
- Gulliford, M. C.; Bicknell, E. J.; Scarpello, J. H., 1989: Differential effect of protein and fat ingestion on blood glucose responses to high- and low-glycemic-index carbohydrates in noninsulin-dependent diabetic subjects. *American Journal of Clinical Nutrition* **50**, 773–777.
- Hannah, S. S., 1999: Role of dietary protein in weight management. *Supplement to Compendium on Continuing Education for the Practicing Veterinarian* **21**, 11.
- Harrison, H. E.; Darrow, D. C.; Yannet, H., 1936: The total electrolyte content of animals and its probable relation to the distribution of body water. *Journal of Biological Chemistry* **113**, 515–529.
- Hewson-Hughes, A. K.; Gilham, M. S.; Upton, S.; Colyer, A.; Butterwick, R.; Miller, A. T., 2011: The effect of dietary starch level on postprandial glucose and insulin concentrations in cats and dogs. *British Journal of Nutrition* **106**, 105–109.
- Hewson-Hughes, A. K.; Hewson-Hughes, V. L.; Colyer, A.; Miller, A. T.; McGrane, S. J.; Hall, S. R.; Butterwick, R. F.; Simpson, S. J.; Raubenheimer, D., 2013: Geometric analysis of macronutrient selection in breeds of the domestic dog, *Canis lupus familiaris*. *Behavioral Ecology* **24**, 293–304.
- Holm, J.; Lundquist, I.; Björck, I.; Eliasson, A. C.; Asp, N. G., 1988: Degree of starch gelatinization, digestion rate of starch in vitro, and metabolic response in rats. *American Journal of Clinical Nutrition* **47**, 1010–1016.
- Jenkins, D. J.; Wolever, T. M.; Taylor, R. H.; Barker, H.; Fielden, H.; Baldwin, J. M.; Bowling, A. C.; Newman, H. C.; Jenkins, A. L.; Goff, D. V., 1981: Glycemic index of foods: a physiological basis for carbohydrate exchange. *American Journal of Clinical Nutrition* **34**, 362–366.
- Kang, M. H.; Kim, D. H.; Jeong, I. S.; Choi, G. C.; Park, H. M., 2015: Evaluation of four portable blood glucose meters in diabetic and non-diabetic dogs and cats. *Veterinary Quarterly* 1–8.
- Kendall, P. T.; Blaza, S. E.; Holme, D. W., 1982: Assessment of endogenous nitrogen output in adult dogs of contrasting size using a protein-free diet. *Journal of Nutrition* **112**, 1281–1286.
- Kendall, C. W.; Augustin, L. S.; Emam, A.; Josse, A. R.; Saxena, N.; Jenkins, D. J., 2006: The glycemic index: methodology and use. *Nestlé Nutrition Workshop Series Clinical Performance Programme* **11**, 43–53.
- Laflamme, D., 1997: Development and validation of a body condition score system for dogs. *Canine Practice* **22**, 10–15.
- Laflamme, D. P., 2006: Understanding and managing obesity in dogs and cats. *Veterinary Clinics of North America: Small Animal Practice* **36**, 1283–1295.
- Laflamme, D. P., 2011: Companion animals symposium: obesity in dogs and cats: what is wrong with being fat? *Journal of Animal Science* **90**, 1653–1662.
- Love, L.; Cline, M. G., 2015: Perioperative physiology and pharmacology in the obese small animal patient. *Veterinary Anaesthesia and Analgesia* **42**, 119–132.
- Mao, J.; Xia, Z.; Chen, J.; Yu, J., 2013: Prevalence and risk factors for canine obesity surveyed in veterinary practices in Beijing, China. *Preventive Veterinary Medicine* **112**, 438–442.
- Mitsuhashi, Y.; Nagaoka, D.; Bigley, K.E.; Umeda, T.; Otsuji, K.; Bauer, J.E., 2012: Metabolic and hormonal alterations with diacylglycerol and low glycemic index starch during canine weight loss. *ISRN Veterinary Science*, **75059**, 3.
- Moore, D. R.; Churchward-Venne, T. A.; Witard, O.; Breen, L.; Burd, N. A.; Tipton, K. D.; Phillips, S. M., 2015: Protein ingestion to stimulate myofibrillar protein synthesis requires greater relative protein intakes in healthy older versus younger men. *Journals of Gerontology Series A: Biological Sciences and Medical Sciences* **70**, 57–62.
- National Research Council, 2006: *Nutrient Requirements of Dogs and Cats*. The National Academies Press, Washington, USA.
- Nguyen, P.; Diez, M., 2010: Obesity. In: S. J. Ettinger, E. C. Feldman (eds), *Textbook of Veterinary Internal Medicine*, 7th ed., Vol. 2. Saunders Elsevier, Canada, pp. 643–652.
- Nguyen, P.; Dumon, H.; Buttin, P.; Martin, L.; Gouro, A. S., 1994: Composition of meal influences changes in postprandial incremental glucose and insulin in healthy dogs. *Journal of Nutrition* **124**, 2707S–2711S.
- Nguyen, P.; Dumon, H.; Biourge, V.; Pouteau, E., 1998: Glycemic and insulino-mic responses after ingestion of commercial foods in healthy dogs: influence of food composition. *Journal of Nutrition* **128**, 2654S–2658S.

- Nuttall, F. Q.; Gannon, M. C., 1990: Metabolic response to egg white and cottage cheese protein in normal subjects. *Metabolism* **39**, 749–755.
- Pourhassan, M.; Bosity-Westphal, A.; Schautz, B.; Braun, W.; Glüer, C. C.; Müller, M. J., 2014: Impact of body composition during weight change on resting energy expenditure and homeostasis model assessment index in overweight nonsmoking adults. *American Journal of Clinical Nutrition* **99**, 779–791.
- R Core Team, 2013: *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>
- Raubenheimer, D.; Machovsky-Capuska, G. E.; Gosby, A. K.; Simpson, S., 2015: Nutritional ecology of obesity: from humans to companion animals. *British Journal of Nutrition* **113**(Suppl), S26–S39.
- Thengchaisri, N.; Theerapun, W.; Kaewmukul, S.; Sastravaha, A., 2014: Abdominal obesity is associated with heart disease in dogs. *BioMed Central Veterinary Research* **10**, 131.
- Verkest, K. R.; Rand, J. S.; Fleeman, L. M.; Morton, J. M., 2012: Spontaneously obese dogs exhibit greater postprandial glucose, triglyceride, and insulin concentrations than lean dogs. *Domestic Animal Endocrinology* **42**, 103–112.
- Weeth, L. P., 2016: Other risks/possible benefits of obesity. *Veterinary Clinics of North America: Small Animal Practice* **46**, 843–853.
- Weijs, P. J.; Wolfe, R. R., 2016: Exploration of the protein requirement during weight loss in obese older adults. *Clinical Nutrition* **35**, 394–398.
- WSAVA Nutritional Assessment Guidelines Task Force Members, Freeman, L.; Becvarova, I.; Cave, N.; MacKay, C.; Nguyen, P.; Rama, B.; Takashima, G.; Tiffin, R.; Tsjimoto, H.; van Beukelen, P., 2011: WSAVA nutritional assessment guidelines. *Journal of Small Animal Practice* **52**, 385–396.

# VETERINARY<sup>TM</sup> HPM


## DIETETIC RANGE

### WEIGHT

# 1

## WEIGHT LOSS & DIABETES

Obesity (>30%) & Diabetes



# VETERINARY<sup>TM</sup> HPM

## DIETETIC RANGE

### WEIGHT

1

### WEIGHT LOSS & DIABETES

Obesity (>30%) & Diabetes

